## (19) World Intellectual Property Organization

International Bureau



## A COLOR SHILLOON HI COLOR COLOR

(43) International Publication Date 2 June 2005 (02.06.2005)

PCT

## (10) International Publication Number WO 2005/049859 A1

(51) International Patent Classification7:

C12Q 1/68

(21) International Application Number:

PCT/EP2003/050844

(22) International Filing Date:

17 November 2003 (17.11.2003)

(25) Filing Language:

English

(26) Publication Language:

English

- (71) Applicant (for all designated States except US): AGI-LENT TECHNOLOGIES, INC. [US/US]; 395 Page Mill Road, Palo Alto 94306 (US).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): LÜBKE, Gerd [DE/DE]; Huelbenstr. 28, 71088 Holzgerlingen (DE).
- (74) Agent: BARTH, Daniel; Herrenberger Str. 130, 71034 Böblingen (DE).


- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

## Published:

with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NUCLEIC ACID DETECTION



(57) Abstract: The invention discloses a method for detection of a target nucleic acid sequence (1A) in a mixture of different nucleic acids (5) having additional binding sites (10) comprising the subsequent steps of A), hybridizing the target nucleic acid sequence with a probe (15) in liquid phase, the probe having a first label (20), separating the different nucleic acids (5) and the target nucleic acid sequence (1 A) in a step B) and in a subsequent step C) detecting the target nucleic acid (1 A) using the labeled probe (15). The method of the invention allows an easy to handle and fast detection of target nucleic acid sequences in nucleic acid mixtures.

